

Subject Code	Q Id	Questions	Answer Key
612	1	<p>A compact subset of \mathbb{R} is</p> <p>(A) $(0,1)$</p> <p>(B) $[0,1)$</p> <p>(C) $(0,1]$</p> <p>(D) $[0,1]$</p>	(D)
612	2	<p>Which of the following function is periodic?</p> <p>(A) $\sin x$</p> <p>(B) e^x</p> <p>(C) $\log x$</p> <p>(D) $\sin^{-1} x$</p>	(A)
612	3	$I = \int_0^a \frac{f(x) dx}{f(x) + f(a-x)} =$ <p>(A) $f(a)$</p> <p>(B) $f(2a)$</p> <p>(C) 0</p> <p>(D) $a/2$</p>	(D)
612	4	<p>If $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, then A^n, for $n \in \mathbb{N}$ is equal to</p> <p>(A) $2^{n-1} A$</p> <p>(B) $2^n A$</p> <p>(C) nA</p>	(A)

		(D) nI	
612	5	<p>If $y = \sqrt{\cos x + \sqrt{\cos x + \sqrt{\cos x + \dots}}}$, then $\frac{dy}{dx}$ is</p> <p>(A) $\frac{\sin x}{1-2y}$</p> <p>(B) $\frac{\sin x}{2y-1}$</p> <p>(C) $\frac{\cos x}{1-2y}$</p> <p>(D) $\frac{\cos x}{2y-1}$</p>	(A)
612	6	<p>For the function $f(z) = \sin(1/z)$, $z = 0$ is</p> <p>(A) an essential singularity</p> <p>(B) a branch point</p> <p>(C) a removable singularity</p> <p>(D) a simple pole</p>	(A)
612	7	<p>The partial differential equation $u_{xx} - xu_{yy} = 0$ is</p> <p>(A) elliptic, $x > 0$</p> <p>(B) hyperbolic, $x < 0$</p> <p>(C) hyperbolic, $x > 0$</p> <p>(D) None of the above</p>	(C)
612	8	<p>The value of $\lim_{x \rightarrow \frac{\pi}{4}} \frac{\int_2^{\sec^2 x} f(t) dt}{x^2 - \frac{\pi^2}{16}}$ is</p> <p>(A) $8 \frac{f(2)}{\pi}$</p> <p>(B)</p>	(A)

		$2\sec^2 x$ (C) $\frac{f(2)}{\pi}$ (D) $f(2)$	
612	9	If $\int_{\sin x}^1 t^2 f(t) dt = 1 - \sin x$, then $f(2)$ is (A) $\frac{1}{9}$ (B) $\frac{1}{4}$ (C) 4 (D) 16	(B)
612	10	The derivative of e^{x^2} with respect to x^4 is (A) $\frac{e^{x^2}}{2x^2}$ (B) $\frac{e^x}{2}$ (C) $\frac{e^{x^2}}{3x^2}$ (D) $\frac{e^{x^2}}{3x}$	(A)
612	11	Consider the functions $f(x) = x \tan x$ and $g(x) = x \cos x^2$. Then (A) both f and g are odd functions (B) f is even function and g is odd (C) both f and g are even functions (D) f is odd function and g is even	(B)
612	12	The domain and range of the function $f(x) = \sin^{-1} x$ are respectively given by	(A)

		<p>(A) $[-1, 1]$ and $(-\infty, \infty)$</p> <p>(B) $[-1, 1]$ and $[0, \pi]$</p> <p>(C) $[-1, 1]$ and $[0, \pi/2]$</p> <p>(D) $[-1, 1]$ and $[\pi, 2\pi]$</p>	
612	13	<p>The value of \sqrt{i} is</p> <p>(A) $1+i$</p> <p>(B) $1-i$</p> <p>(C) -1</p> <p>(D) $\pm \frac{1+i}{\sqrt{2}}$</p>	(D)
612	14	<p>If $x = e^t, y = \sin t, z = \cos t$ and $u = \log(x+y+z)^2$, then $\frac{du}{dt}$ is</p> <p>(A) $\frac{2(e^t + \cos t - \sin t)}{(e^t + \cos t + \sin t)}$</p> <p>(B) $\frac{(e^t + \cos t - \sin t)}{2(e^t + \cos t + \sin t)}$</p> <p>(C) $\frac{2(e^t + \cos t + \sin t)}{(e^t + \cos t - \sin t)}$</p> <p>(D) $\frac{2(e^t - \cos t + \sin t)}{(e^t + \cos t - \sin t)}$</p>	(A)
612	15	<p>The value of $\left((i^{999})^2 + \left(\frac{1}{i^{-2003}} \right)^2 \right)^2$ is</p> <p>(A) -4</p> <p>(B) 4</p> <p>(C) -1</p>	(B)

		(D) 1	
612	16	<p>The intersection of $\{x: x^2 - 26 \leq 10\}$ and $\{x: x^2 - 5 \leq 4\}$</p> <p>(A) is a closed interval</p> <p>(B) is an open interval</p> <p>(C) is empty</p> <p>(D) contains exactly 4 points</p>	(C)
612	17	<p>If in a group, an element a has order 65, then the order of a^{25} is</p> <p>(A) 5</p> <p>(B) 12</p> <p>(C) 13</p> <p>(D) 7</p>	(C)
612	18	<p>A harmonic conjugate $v(x, y)$ of the function $u(x, y) = x^3 - 3xy^2$ on \mathbb{C} is</p> <p>(A) $y^3 - 3x^3 + c$</p> <p>(B) $3x^2y - y^3 + c$</p> <p>(C) $x^3 - 3x^2y + c$</p> <p>(D) $y^2 - 3xy + c$</p>	(B)
612	19	<p>If $\left(\frac{1-i}{1+i}\right)^{100} = a + ib$, then</p> <p>(A) $a = 2, b = -1$</p> <p>(B) $a = 1, b = 0$</p> <p>(C) $a = 0, b = 1$</p> <p>(D) $a = -1, b = 2$</p>	(B)
612	20		(C)

		<p>Which of the following functions is continuous but not differentiable?</p> <p>(A) x</p> <p>(B) e^x</p> <p>(C) x</p> <p>(D) $\sin x$</p>	
612	21	<p>In which of the following subspaces, the sequence $\left\{\frac{1}{n}\right\}$ is Cauchy but not convergent?</p> <p>(A) $[0,1]$</p> <p>(B) $[0,1)$</p> <p>(C) $(0,1]$</p> <p>(D) $(0,1)$</p>	(C)
612	22	<p>If $f(x) = 1 + \frac{\log x}{1!} + \frac{(\log x)^2}{2!} + \dots$, then $\int f(x) dx$ is</p> <p>(A) $\log x + c$</p> <p>(B) $e^x + c$</p> <p>(C) $x + c$</p> <p>(D) $\frac{x^2}{2} + c$</p>	(D)
612	23	<p>If $F(1) = 2$ and $F(n) = F(n-1) + \frac{1}{2}$ for all integers $n > 1$, then $F(51) =$</p> <p>(A) 25</p> <p>(B) 26</p> <p>(C) 27</p> <p>(D) 28</p>	(C)
612	24		(C)

		<p>The value of $\int_C \frac{e^z dz}{z^4}$ when C is $z =1$ is</p> <p>(A) $\frac{8\pi i}{3}$</p> <p>(B) $\frac{4\pi i}{3}$</p> <p>(C) $\frac{\pi i}{3}$</p> <p>(D) $\frac{2\pi i}{3}$</p>	
612	25	<p>For each x in $[0, 1]$, let $f(x) = x$ if x is rational and $f(x) = 1 - x$ if x is irrational. Then</p> <p>(A) $f(x+1) = f(x)$</p> <p>(B) $f(x) - f(1-x) = 1$</p> <p>(C) $f(1-x) - f(x) = 1$</p> <p>(D) $f(x) + f(1-x) = 1$</p>	(D)
612	26	<p>The area under one arc of the cycloid $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$</p> <p>(A) $\frac{\pi a^2}{8}$</p> <p>(B) $\frac{3\pi a^2}{16}$</p> <p>(C) $3\pi a^2$</p> <p>(D) $\frac{3\pi a^2}{32}$</p>	(C)
612	27	<p>If A is a self adjoint matrix, then its diagonal entries are</p> <p>(A) all complex numbers</p> <p>(B) all real numbers</p> <p>(C) 0</p>	(B)

		(D) - 1	
612	28	<p>The differential equation obtained by eliminating f from $z = f(x^2 + y^2)$ when $p = \frac{\partial z}{\partial x}$ and $q = \frac{\partial z}{\partial y}$ is</p> <p>(A) $py = qx$</p> <p>(B) $pq = xy$</p> <p>(C) $px = qy$</p> <p>(D) $x = y$</p>	(A)
612	29	<p>The differential equation of the family of curves $y = e^{2x}(A \cos x + B \sin x)$ where A and B are constants is</p> <p>(A) $\frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} + 4y = 0$</p> <p>(B) $\frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} + 5y = 0$</p> <p>(C) $\frac{d^2 y}{dx^2} - 4y \frac{dy}{dx} + 5y = 0$</p> <p>(D) $\frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} + 4x = 0$</p>	(B)
612	30	<p>The value of $\lim_{x \rightarrow \infty} \int_0^x e^{-t^2} dt$ is</p> <p>(A) $\sqrt{\pi}$</p> <p>(B) $\sqrt{\frac{\pi}{2}}$</p> <p>(C) $\frac{\sqrt{\pi}}{2}$</p> <p>(D) π</p>	(C)
612	31	<p>If $u = \tan^{-1}\left(\frac{y}{x}\right)$, then $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$ is equal to</p> <p>(A)</p>	(C)

		$\frac{2xy}{x^2 + y^2}$ <p>(B)</p> 1 <p>(C)</p> 0 <p>(D)</p> $\frac{x^2}{x^2 + y^2}$	
612	32	<p>The residue of $\frac{z^2}{(z-1)(z-2)(z-3)}$ at $z=1$ is</p> <p>(A)</p> -8 <p>(B)</p> $1/2$ <p>(C)</p> -6 <p>(D)</p> 0	(B)
612	33	<p>If R is a commutative ring and $N = \{x \in R : x^n = 0 \text{ for some integer } n\}$, then</p> <p>(A)</p> $N \text{ is an ideal of } R$ <p>(B)</p> $N \text{ is not an ideal of } R$ <p>(C)</p> $N \text{ is a subring of } R$ <p>(D)</p> $N \text{ is a subfield of } R$	(A)
612	34	<p>The common area in square units between the curves $y^2 = 4x$ and $x^2 = 4y$ is</p> <p>(A)</p> $\frac{16}{5}$ <p>(B)</p> $\frac{16}{3}$ <p>(C)</p> $\frac{8}{3}$ <p>(D)</p> $\frac{8}{5}$	(B)
612	35		(C)

		<p>The value of $(1+i)(1+i^2)(1+i^3)(1+i^4)\dots(1+i^n)$ for $n > 1$ is</p> <p>(A) 1</p> <p>(B) -1</p> <p>(C) 0</p> <p>(D) i</p>	
612	36	<p>The function $f(x) = \frac{x^2 - x}{x}$ is</p> <p>(A) continuous everywhere</p> <p>(B) continuous except at $x = 0$</p> <p>(C) continuous at $x = 0$</p> <p>(D) discontinuous everywhere</p>	(A)
612	37	<p>$\frac{d}{dx} \int_{\sin^2 x}^{2\sin x} e^{t^2} dt$ at $x = \pi$ is</p> <p>(A) 1</p> <p>(B) -1</p> <p>(C) 2</p> <p>(D) -2</p>	(D)
612	38	<p>A ring in which the nonzero elements form a group is called</p> <p>(A) an integral domain</p> <p>(B) a skew-field</p> <p>(C) a field</p> <p>(D) commutative ring</p>	(B)
612	39	<p>\mathbb{R}^n is not a field when</p> <p>(A) $n = 1$</p> <p>(B)</p>	(C)

		$n = 2$ (C) $n > 1$ (D) n is prime	
612	40	The period of the function $f(x) = 3 - 2\cos^2\left(\frac{\pi x}{3}\right)$ (A) 2 (B) 3 (C) 5 (D) 6	(B)
612	41	If $f(x) = \int_0^x e^{-t^2} dt$, then $f'(x)$ is (A) 0 (B) e^{-x^2} (C) e^{-x^2} (D) $e^{-x^2} + c$	(C)
612	42	Suppose f is continuous on $[a, b]$, differentiable on (a, b) and $f(a) = f(b)$. Then for some $c \in (a, b)$, we have (A) $f(a) = f(c)$ (B) $f(b) = f(c)$ (C) $f(c) = \frac{f(b) - f(a)}{b - a}$ (D) $f'(c) = 0$	(D)
612	43	The sum of all the external forces on a system of particles is zero. Which of the following must be true of the system? (A) The total mechanical energy is constant (B)	(B)

		<p>The total linear momentum is constant</p> <p>(C)</p> <p>The total kinetic energy is constant</p> <p>(D)</p> <p>The total potential energy is constant</p>	
612	44	<p>In a field a bull is grazing around a tree in an elliptical path having the tree at a focus. The shortest and the longest distance from the bull and the tree are 200 and 500 units. The eccentricity of the path is</p> <p>(A)</p> $\frac{3}{7}$ <p>(B)</p> $\frac{7}{3}$ <p>(C)</p> $\frac{1}{3}$ <p>(D)</p> $\frac{5}{7}$	(A)
612	45	<p>The solution of the boundary value problem $y'' + 4y = 0, y(0) = 1, y(\pi) = 1$ is</p> <p>(A)</p> $y = \cos 2x$ <p>(B)</p> $y = 0$ <p>(C)</p> $y = \cos 2x + A \sin 2x$, where A is arbitray <p>(D)</p> $y = \sin 2x$	(C)
612	46	<p>Which of the following surface intersects the plane $x = 2$ at a parabola?</p> <p>(A)</p> $-\frac{z^2}{2} = \frac{x^2}{9} + \frac{y^2}{4}$ <p>(B)</p> $\frac{z}{4} = \frac{x}{4} + \frac{y}{9} - 1$ <p>(C)</p> $\frac{z}{2} = \frac{x^2}{9} - \frac{y^2}{4}$ <p>(D)</p> $\frac{z^2}{4} = \frac{x^2}{9} + \frac{y^2}{4} + 1$	(C)
612	47	<p>An integer solution of $(1-i)^x = 2^{\frac{x}{2}}$ is given by</p> <p>(A)</p> 8	(A)

		<p>(B) 1</p> <p>(C) 6</p> <p>(D) 2</p>	
612	48	<p>Let P be the point (1,0) and let Q be a point on the locus $y^2 = 4x$. The locus of mid point of PQ is</p> <p>(A) $y^2 + 2x + 1 = 0$</p> <p>(B) $y^2 - 2x + 1 = 0$</p> <p>(C) $x^2 - 2y + 1 = 0$</p> <p>(D) $x^2 + 2y + 1 = 0$</p>	(B)
612	49	<p>The angle between the lines $6x = 3y = 4z$ and $2x = -y = z$</p> <p>(A) $\frac{\pi}{3}$</p> <p>(B) 0</p> <p>(C) $\frac{\pi}{4}$</p> <p>(D) $\frac{\pi}{2}$</p>	(D)
612	50	<p>If α, β are the roots of the equation $x^2 - 2x + 4 = 0$, then $\alpha^6 + \beta^6$ is equal to</p> <p>(A) 64</p> <p>(B) -64</p> <p>(C) 0</p> <p>(D) -128</p>	(C)
612	51	<p>Let $f(x) \in R[x] = \{a_0 + a_1x + \dots + a_nx^n : a_i \in R, \text{ a ring and } n \text{ is a non-negative integer}\}$. If $f(a) = f'(a) = 0$, then $(x-a)^2$ divides</p> <p>(A) $f(x)$</p>	(A)

		<p>(B) $f'(x)$</p> <p>(C) $f(x) - a$</p> <p>(D) $f'(x) - a$</p>	
612	52	<p>If a group G is such that $(a \cdot b)^2 = a^2 \cdot b^2$, then G is</p> <p>(A) non-abelian</p> <p>(B) abelian</p> <p>(C) cyclic</p> <p>(D) non-cyclic</p>	(B)
612	53	<p>If $f(x) = 2x^3 + x^2 + 2x + 2$ and $g(x) = 2x^2 + 2x + 1$ are in $Z_3[x]$, then $f(x) + g(x)$ is</p> <p>(A) $2x^3 + 3x^2 + 4x + 3$</p> <p>(B) $2x^3 + x$</p> <p>(C) $3x^3 + 4x + 3$</p> <p>(D) $2x^3 + x^2 + 2x + 2$</p>	(B)
612	54	<p>If three forces acting at a point are in equilibrium, then they can be represented in magnitude and direction by</p> <p>(A) the medians of a triangle</p> <p>(B) the sides of a triangle</p> <p>(C) the altitudes of a triangle</p> <p>(D) the perpendicular bisectors of a triangle</p>	(B)
612	55	<p>Let G be group of order 49. Then</p> <p>(A) G is abelian</p> <p>(B) G is cyclic</p> <p>(C)</p>	(A)

		<p>G is non-abelian</p> <p>(D) centre of G has order 7</p>	
612	56	<p>Let $\lambda = e^{\frac{5\pi i}{6}}$. Then the smallest positive integer n such that $\lambda^n = 1$ is</p> <p>(A) 6</p> <p>(B) 9</p> <p>(C) 12</p> <p>(D) 5</p>	(C)
612	57	<p>If $f(x)$ is an odd function and $g(x)$ is an even function, then</p> <p>(A) $f \circ g$ is odd</p> <p>(B) $f \circ g$ is even</p> <p>(C) $f \circ f$ is even</p> <p>(D) $g \circ g$ is odd</p>	(B)
612	58	<p>Let $f(z) = \sin\left(\frac{1}{1-z}\right)$. Then $z=1$ is a</p> <p>(A) non-isolated essential singularity</p> <p>(B) removable singularity</p> <p>(C) isolated essential singularity</p> <p>(D) analytic</p>	(C)
612	59	<p>Let P and Q be square matrices such that $PQ = I$, the identity matrix. Then zero is an eigen value of</p> <p>(A) P but not of Q</p> <p>(B) Q but not of P</p> <p>(C) Both P and Q</p>	(D)

		(D) Neither P nor Q	
612	60	<p>A root of the equation $e^x = 4x$ lies between</p> <p>(A) – 1 and 0</p> <p>(B) 1 and 2</p> <p>(C) 1 and 3</p> <p>(D) 3 and 4</p>	(C)
612	61	<p>The value of $\nabla^2 \log r$ is equal to</p> <p>(A) $\frac{1}{r}$</p> <p>(B) $\frac{1}{r^2}$</p> <p>(C) $-\frac{1}{r^2}$</p> <p>(D) r^2</p>	(C)
612	62	<p>$\bigcup_{n=1}^{\infty} \left[1 - \frac{1}{n}, 1 + \frac{1}{n} \right] =$</p> <p>(A) $[0, 2]$</p> <p>(B) $[0, 1]$</p> <p>(C) $[1, 2]$</p> <p>(D) $\{1\}$</p>	(A)
612	63	<p>If $x = e^{y+e^{y+x}}$, then $\frac{dy}{dx}$ is</p> <p>(A) $\frac{x(1-x)}{y}$</p> <p>(B) $\frac{y}{x(1-x)}$</p>	(C)

		(C) $\frac{1-x}{x}$ (D) e^{x+y}	
612	64	The value of $\int_C \frac{\sin z}{z} dz$, when C is $ z =1$, is (A) 1 (B) π (C) 0 (D) -1	(C)
612	65	The residue of $\frac{1}{(z^2+1)^3}$ at $z=i$ is (A) $\frac{5}{16i}$ (B) $\frac{3}{13i}$ (C) $\frac{3}{16i}$ (D) $\frac{5}{13i}$	(C)
612	66	The inverse transform of $T(z) = \frac{z+2}{z+3}$ is (A) $\frac{2-3\omega}{\omega-1}$ (B) $\frac{2+3\omega}{\omega-1}$ (C) $\frac{2-3\omega}{\omega+1}$ (D) $\frac{2+3\omega}{\omega+1}$	(A)
612	67		(B)

		<p>Which of the following equations is exact?</p> <p>(A)</p> $\frac{dy}{dx} = \frac{(x^2 - 2xy + 3y^2)}{(y^2 + 6xy - x^2)}$ <p>(B)</p> $\frac{dy}{dx} = -\frac{(x^2 - 2xy + 3y^2)}{(y^2 + 6xy - x^2)}$ <p>(C)</p> $\frac{dy}{dx} = \frac{(e^y \sin x)}{e^y (\sin x + 1)}$ <p>(D)</p> $\frac{dy}{dx} = -\frac{(e^y \sin x)}{e^y (\sin x + 1)}$	
612	68	<p>The unit normal to the surface $x^2 + y^2 + z^2 = 6$ at the point $(2, 1, 1)$ is</p> <p>(A)</p> $2\vec{i} - \vec{j} + \vec{k}$ <p>(B)</p> $\frac{1}{\sqrt{6}}(2\vec{i} + \vec{j} + \vec{k})$ <p>(C)</p> $\frac{1}{6}(2\vec{i} + \vec{j} + \vec{k})$ <p>(D)</p> $\frac{1}{\sqrt{6}}(2\vec{i} - \vec{j} + \vec{k})$	(B)
612	69	<p>The solution of $\frac{d^2y}{dx^2} = 1 + y$ which vanishes at $x = 0$ and tends to a finite limit as $x \rightarrow \infty$ is</p> <p>(A)</p> $(1 + e^{-x})$ <p>(B)</p> $(e^{-x} - 1)$ <p>(C)</p> $(e^x + e^{-x} - 2)$ <p>(D)</p> $(e^x - 1)$	(B)
612	70	<p>If $1, \omega, \omega^2$ are cube roots of unity, then $(1 + \omega)^3 - (1 + \omega^2)^3$ is</p> <p>(A)</p> <p>1</p> <p>(B)</p>	(C)

		<p>-1</p> <p>(C)</p> <p>0</p> <p>(D)</p> <p>2</p>	
612	71	<p>The solution of the differential equation $x \frac{dy}{dx} + y = x^2$ is</p> <p>(A)</p> $3y = x^2 + \frac{c}{x}$ <p>(B)</p> $y = \frac{x^3}{3} + c$ <p>(C)</p> $x^2 y = \frac{x^3}{3} + c$ <p>(D)</p> $xy = c$	(A)
612	72	<p>The set $[0,1]$ is</p> <p>(A)</p> <p>countable</p> <p>(B)</p> <p>countably finite</p> <p>(C)</p> <p>uncountable</p> <p>(D)</p> <p>countably infinite</p>	(C)
612	73	<p>The function $f(x) = \begin{cases} 1 & x \text{ is rational} \\ -1 & x \text{ is reational} \end{cases}$ is</p> <p>(A)</p> <p>continuous at $x = 0$</p> <p>(B)</p> <p>discontinuous everywhere</p> <p>(C)</p> <p>discontinuous only at $x = 0$</p> <p>(D)</p> <p>continuous at $x = \frac{1}{2}$</p>	(B)
612	74	<p>The point $z = 0$ of $f(z) = \frac{z+3}{z^2(z^2+2)}$ is</p> <p>(A)</p> <p>an isolated singularity</p>	(A)

		<p>(B) a removable singularity</p> <p>(C) an essential singularity</p> <p>(D) None of the above</p>	
612	75	<p>The solution of $zp + x = 0$ is</p> <p>(A) $x^2 + z^2 = \phi(y)$</p> <p>(B) $x^2 z^2 = \phi(y)$</p> <p>(C) $(x - y)^2 = \phi(y)$</p> <p>(D) $(x + y)^2 = \phi(y)$</p>	(A)
612	76	<p>The solution of the initial value problem $x \frac{dy}{dx} = 3y$, $y(1) = 3$ is</p> <p>(A) $y = x^3$</p> <p>(B) $y^3 = x$</p> <p>(C) $y = 3x^3$</p> <p>(D) $3y^3 = x$</p>	(C)
612	77	<p>If \vec{a} and \vec{b} are two unit vectors and θ is the angle between them, then $(\vec{a} - \vec{b})$ is a unit vector if</p> <p>(A) $\theta = \pi/6$</p> <p>(B) $\theta = \pi/4$</p> <p>(C) $\theta = \pi/3$</p> <p>(D) $\theta = 2\pi/3$</p>	(C)
612	78	<p>Which of the following is the order of a non-abelian group?</p> <p>(A) 4</p> <p>(B) 8</p>	(B)

		<p>(C) 9</p> <p>(D) 13</p>	
612	79	<p>Let S_4 be the group of permutations on four letters. The number of elements of order 2 in the group S_4 is</p> <p>(A) 6</p> <p>(B) 9</p> <p>(C) 4</p> <p>(D) 12</p>	(B)
612	80	<p>If the entries of a 2×2 matrix A are defined by the formula $a_{ij} = i^2 + j^2$, then A is</p> <p>(A) a symmetric matrix</p> <p>(B) a skew symmetric matrix</p> <p>(C) the identity matrix</p> <p>(D) $\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$</p>	(A)
612	81	<p>Which of the following formula is false?</p> <p>(A) $z_1 z_2 = z_1 z_2$</p> <p>(B) $\arg z_1 z_2 = \arg z_1 + \arg z_2 \pmod{2\pi}$</p> <p>(C) $z_1 + z_2 \leq z_1 + z_2$</p> <p>(D) $\arg(z_1 + z_2) = \arg z_1 + \arg z_2 \pmod{2\pi}$</p>	(D)
612	82	<p>Which of the following functions $f: R \rightarrow R$ is one-one and onto?</p> <p>(A) $f(x) = x^3 + 2$</p> <p>(B) $f(x) = \sin x$</p> <p>(C) $f(x) = \cos x$</p>	(A)

		(D) $f(x) = x^4 - x^2$	
612	83	If $I_n = \left(-\frac{1}{n}, \frac{1}{n}\right)$, $n \in \mathbb{N}$, then $\bigcap_{n=1}^{\infty} I_n$ is (A) open (B) not open (C) neither open nor closed (D) unbounded set	(B)
612	84	If $I = \int_0^{\pi/2} \sin^8 x dx$ and $J = \int_0^{\pi/2} \cos^8 x dx$, then the value of I/J is (A) $5\pi/16$ (B) $5\pi/32$ (C) π (D) 1	(D)
612	85	The Laplace transform of $\frac{\sin at}{at}$ is (A) $\tan\left(\frac{a}{s}\right)$ (B) $\tan^{-1}\left(\frac{a}{s}\right)$ (C) $\frac{\pi}{2} - \tan^{-1}\left(\frac{a}{s}\right)$ (D) $\frac{\pi}{2} - \tan\left(\frac{s}{a}\right)$	(C)
612	86	The value of $\int_{-1}^{+1} x \ln x $ is (A) -1 (B) 1 (C)	(B)

		0 (D) 2	
612	87	<p>Let there be three distinct prime numbers C, A, T such that the product $C \times A \times T = 2014$. The possible value of the sum $C + A + T$ is</p> <p>(A) 58</p> <p>(B) 74</p> <p>(C) 109</p> <p>(D) 214</p>	(B)
612	88	<p>a, b, c ($a > c$) are three digits, from left to right, of a three digit number. If the number with these digits reversed is subtracted from the original number, the resulting number has digit 4 in its unit place. The other two digits from left to right are</p> <p>(A) 5 and 4</p> <p>(B) 5 and 9</p> <p>(C) 4 and 5</p> <p>(D) 9 and 5</p>	(B)
612	89	<p>If 3^{2015} is divided by 11, then the remainder is</p> <p>(A) 0</p> <p>(B) 1</p> <p>(C) 4</p> <p>(D) 5</p>	(B)
612	90	<p>If 12 divides $ab313ab$, the smallest value of $a + b$ is</p> <p>(A) 2</p> <p>(B) 4</p> <p>(C) 6</p> <p>(D) 7</p>	(B)

612	91	<p>The sum to infinity of the G.P. $-\frac{5}{4}, \frac{5}{16}, -\frac{5}{64}, \dots$ is</p> <p>(A) -1</p> <p>(B) 0</p> <p>(C) $\frac{25}{256}$</p> <p>(D) $-\infty$</p>	(A)
612	92	<p>The cube roots of 1 are in</p> <p>(A) A.P.</p> <p>(B) G.P.</p> <p>(C) H.P</p> <p>(D) A.P. but not H.P.</p>	(B)
612	93	<p>The greatest number of three digits which when added to 45 is exactly divisible by 6, 8, 12 is</p> <p>(A) 963</p> <p>(B) 987</p> <p>(C) 999</p> <p>(D) 1044</p>	(B)
612	94	<p>The cube roots of 8 are</p> <p>(A) $2, -2$</p> <p>(B) $2, 2i, 2i^2$</p> <p>(C) $2, -2, 2i$</p> <p>(D) $2, 2\omega, 2\omega^2$</p>	(D)
612	95	<p>The amplitude of $(-1)^5$ is</p> <p>(A) $\pi/2$</p> <p>(B)</p>	(D)

		$\pi/4$ (C) $3\pi/2$ (D) π	
612	96	The principle value of the amplitude of $-1 + i\sqrt{3}$ is (A) $\pi/2$ (B) $\pi/3$ (C) $2\pi/3$ (D) π	(C)
612	97	If $1, \omega, \omega^2$ are three roots of unity, then $(3 + \omega^2 + \omega^4)^6$ is (A) 0 (B) 2 (C) 64 (D) 729	(C)
612	98	The complex number $1 + i$ in the polar form is (A) $(2, \pi/2)$ (B) $(\sqrt{2}, \pi/4)$ (C) $(3, 2\pi/3)$ (D) $(\sqrt{3}, \pi/4)$	(B)
612	99	The value of $(1 + i)^6 + (1 - i)^6$ is (A) 32 (B) 8 (C) 1 (D) 0	(D)
612	100		(A)

		<p>The coefficient of x^{99} in the expansion $(x-1)(x-2)\dots(x-100)$ is</p> <p>(A) -5050</p> <p>(B) -4950</p> <p>(C) -4851</p> <p>(D) -100</p>	
612	101	<p>The imaginary part of $\sin(x+iy)$ is</p> <p>(A) $\sin x \cos hy$</p> <p>(B) $i \sin x \sin hy$</p> <p>(C) $\cos x \cos hy$</p> <p>(D) $\cos x \sin hy$</p>	(D)
612	102	<p>If $a > 0, b > 0, c > 0$, then the correct symbol in to the following $a^2 + b^2 + c^2$ $ab + bc + ca$ is</p> <p>(A) $<$</p> <p>(B) $>$</p> <p>(C) \leq</p> <p>(D) \geq</p>	(D)
612	103	<p>The diagonal of a cube is $\sqrt{12}$ cm. The volume of the cube in cubic cms. is</p> <p>(A) $3\sqrt{2}$</p> <p>(B) $12\sqrt{12}$</p> <p>(C) 24</p> <p>(D) 8</p>	(D)
612	104	<p>If $a-b=3$ and $a^3-b^3=117$, then $a+b$ is equal to</p> <p>(A) 39</p> <p>(B) 29</p>	(D)

		(C) $\frac{9}{7}$	
612	105	If $x + 1/x = 3$, then $x^4 + 1/x^4$ is equal to (A) 279 (B) 168 (C) 81 (D) 47	(D)
612	106	The maximum value of $4 - x - x^2$ is (A) 10 (B) 4 (C) $\frac{21}{4}$ (D) $\frac{17}{4}$	(D)
612	107	The value of $\log_3 \log_2 \log_{\sqrt{6}} 81$ is equal to (A) 1 (B) 7 (C) 11 (D) 22	(A)
612	108	If $x = \log 3/4$, $y = \log 4/7$, $z = 2 \log \sqrt[4]{(7/4)}$, then the value of $13^{2(x+y+z)}$ is (A) 1 (B) 7 (C) 11 (D) 22	(A)

612	109	<p>The number of solutions of the equation $x^{\log_{10} x} = 100x$ is</p> <p>(A) 0</p> <p>(B) 1</p> <p>(C) 2</p> <p>(D) 3</p>	(C)
612	110	<p>The value of $\cot 9^\circ \cot 27^\circ \cot 63^\circ \cot 81^\circ$ is</p> <p>(A) 1</p> <p>(B) $\sqrt{2}/3$</p> <p>(C) $1/\sqrt{2}$</p> <p>(D) 1/2</p>	(A)
612	111	<p>If $\sin 2\theta = \cos \theta$, then the value of θ is</p> <p>(A) $\pi/2$</p> <p>(B) $\pi/4$</p> <p>(C) $3\pi/2$</p> <p>(D) π</p>	(C)
612	112	<p>The value of $\frac{\sin 2x}{1 + \cos 2x}$ is equal to</p> <p>(A) $\sin x$</p> <p>(B) $\cos x$</p> <p>(C) $\tan x$</p> <p>(D) $\cot x$</p>	(C)
612	113	<p>If $\sin x \cos y = 1/4$ and $3 \tan x = \tan y$, then the value of $\sin(x + y)$ is equal to</p> <p>(A) 0</p> <p>(B) 1</p>	(B)

		(C) 3 (D) 4	
612	114	<p>ABC is a right-angled triangle with $\angle B = 90^\circ$. M is the midpoint of AC and $BM = \sqrt{117} \text{ cm}$. Sum of the other two sides AB and BC is 30cm. The area of the triangle in sq.cms. is</p> (A) 27 (B) 108 (C) 110 (D) 112	(B)
612	115	<p>The area of the region in the Cartesian plane whose points (x, y) satisfy $x + y + x+y \leq 2$ is</p> (A) 2 (B) 2.5 (C) 3 (D) 4	(C)
612	116	<p>The equation of tangent to the circle $x^2 + y^2 = 85$ at the point (7, 6) is</p> (A) $7x - 6y = 85$ (B) $6x + 7y = \sqrt{85}$ (C) $6x - 7y = \sqrt{85}$ (D) $7x + 6y = 85$	(D)
612	117	<p>The equation of the normal of $y^2 = 20x$ at (5, 10) is</p> (A) $7x - 6y = 85$ (B) $6x + 7y = \sqrt{85}$ (C) $6x - 7y = \sqrt{85}$ (D) $7x + 6y = 85$	(B)

612	118	<p>The centre of an ellipse $9x^2 + 5y^2 - 36x - 50y - 164 = 0$ is at</p> <p>(A) (2,5)</p> <p>(B) (1,-2)</p> <p>(C) (-2,1)</p> <p>(D) (0,0)</p>	(A)
612	119	<p>If $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, then $(A - A^t)$, [where A^t is transpose of matrix A], is</p> <p>(A) null matrix</p> <p>(B) identity matrix</p> <p>(C) symmetric</p> <p>(D) skew-symmetric</p>	(D)
612	120	<p>The value of the determinant $\begin{vmatrix} x & 1 & y+z \\ y & 1 & z+x \\ z & 1 & x+y \end{vmatrix}$</p> <p>(A) $x+y+z$</p> <p>(B) 0</p> <p>(C) 1</p> <p>(D) $1+x+y+z$</p>	(B)
612	121	<p>Let M and N be two non-empty subsets of a set X such that M is not a subset of N. Then</p> <p>(A) N is a subset of M</p> <p>(B) M and the complement of N are non-disjoint</p> <p>(C) M and N are disjoint</p> <p>(D) M is not a subset of the complement of N</p>	(B)
612	122	<p>If $X = \{ 2, 4, 6, 8 \}$ and $Y = \{ 2, 3, 4, 5 \}$ be two sets, then the symmetric difference of X and Y is the set</p> <p>(A) $\{ 6, 8 \}$</p>	(D)

		<p>(B) $\{3, 5\}$</p> <p>(C) $\{2, 4, 3, 5\}$</p> <p>(D) $\{3, 5, 6, 8\}$</p>	
612	123	<p>Out of 84 students, the number of students taking Mathematics is 65 and the number of students taking both Mathematics and Statistics is 20. Then the number of students taking only Statistics is</p> <p>(A) 19</p> <p>(B) 39</p> <p>(C) 45</p> <p>(D) 64</p>	(B)
612	124	<p>The number of integers between 1 and 6300 inclusive which are divisible neither by 5 nor by 3 is</p> <p>(A) 5040</p> <p>(B) 3360</p> <p>(C) 2100</p> <p>(D) 1260</p>	(B)
612	125	<p>A book-shelf holds 5 different computer books, 6 different statistics books and 10 different mathematics books. The number of ways selecting 2 books, one in two subjects is</p> <p>(A) 300</p> <p>(B) 260</p> <p>(C) 140</p> <p>(D) 60</p>	(C)
612	126	<p>Which of the properties is not satisfied for the relation defined by $R = \{ (a, b) : a, b \in Z, \text{ the set of integers, } a - b \leq 3 \}$.</p> <p>(A) reflexive</p> <p>(B) symmetric</p> <p>(C) transitive</p>	(C)

		(D) symmetric and transitive	
612	127	<p>If the functions f and g are defined by $f = \{(5,2), (6,3)\}$ and $g = \{(2,5), (3,6)\}$, then the value of $(f \circ g)(2)$ is</p> <p>(A) 2</p> <p>(B) 3</p> <p>(C) 5</p> <p>(D) 6</p>	(A)
612	128	<p>Let $f(x)$ be a polynomial of degree 1. If $f(10) - f(5) = 15$, then $f(20) - f(5)$ is</p> <p>(A) 25</p> <p>(B) 40</p> <p>(C) 45</p> <p>(D) 65</p>	(C)
612	129	<p>If $\frac{{}^nP_5}{{}^nP_3} = 20$, then value of $n =$</p> <p>(A) 2</p> <p>(B) 8</p> <p>(C) 12</p> <p>(D) 15</p>	(B)
612	130	<p>${}^nC_1 + {}^nC_2 + {}^nC_3 + \dots + {}^nC_n =$</p> <p>(A) $2^n C_n$</p> <p>(B) ${}^{n+1}C_n$</p> <p>(C) 2^n</p> <p>(D) $2^n - 1$</p>	(D)
612	131	<p>${}^{50}C_{11} + {}^{50}C_{12} + {}^{51}C_{13} - {}^{52}C_{13} =$</p>	(D)

		<p>(A) ${}^{52}C_{14}$</p> <p>(B) ${}^{53}C_{13}$</p> <p>(C) $2^{53}C_{12}$</p> <p>(D) 0</p>	
612	132	<p>If $f(x) = x^3 + ax + 1$ and $f(1) = 1$, then the value of $f(2)$ is</p> <p>(A) 1</p> <p>(B) 3</p> <p>(C) 5</p> <p>(D) 7</p>	(D)
612	133	<p>If $f(x) = \frac{x-1}{x+1}$, then $f\left(-\frac{1}{x}\right)$ is equal to</p> <p>(A) $-x$</p> <p>(B) $-1/x$</p> <p>(C) $-f'(x)$</p> <p>(D) $-f\left(\frac{1}{x}\right)$</p>	(D)
612	134	<p>If $f(x) = \frac{x+2}{x-2}$, then $\lim_{x \rightarrow \infty} f(x)$ is</p> <p>(A) 0</p> <p>(B) 1</p> <p>(C) 2</p> <p>(D) does not exist</p>	(B)
612	135	<p>The function $f(x)$ has a minimum at a point $x = a$ if</p> <p>(A) $f'(a) = 0$ and $f''(a) = 0$</p> <p>(B) $f'(a) = 0$ and $f''(a) < 0$</p>	(C)

		<p>(C) $f'(a) = 0$ and $f''(a) > 0$</p> <p>(D) $f'(a) = 0$ and $f''(a) \leq 0$</p>	
612	136	<p>The value of $\int_a^b \frac{1}{(1+x^2)} dx$ is</p> <p>(A) $b^2 - a^2$</p> <p>(B) $\sin^{-1} b - \sin^{-1} a$</p> <p>(C) $\cos^{-1} b - \cos^{-1} a$</p> <p>(D) $\tan^{-1} b - \tan^{-1} a$</p>	(D)
612	137	<p>The value of $\int_{-n}^n a da$ is</p> <p>(A) n^2</p> <p>(B) $n^2 / 4$</p> <p>(C) 0</p> <p>(D) $-a^2$</p>	(A)
612	138	<p>$\int \frac{\log x}{x}$ is equal to</p> <p>(A) $\log(\log x)$</p> <p>(B) $(\log x)^2$</p> <p>(C) $\frac{1}{2} (\log x)^2$</p> <p>(D) $\frac{1}{2} \log(\log)$</p>	(C)
612	139	<p>The equation $y^2 - x^2 + 2x - 1 = 0$ represents</p> <p>(A) a pair of straight lines</p> <p>(B) a circle</p> <p>(C) an ellipse</p> <p>(D) a hyperbola</p>	(A)
612	140		(A)

		<p>The probability of Mr. A solving a problem is $\frac{3}{4}$ and Mr. B solving the same problem is $\frac{3}{4}$. The probability that the problem is solved is</p> <p>(A) 1</p> <p>(B) $\frac{1}{2}$</p> <p>(C) $\frac{3}{16}$</p> <p>(D) $\frac{13}{16}$</p>	
612	141	<p>In a Cricket test series of five tests, the captain of Indian team decides to call heads at every toss. The probability of his winning the toss in all the tests is</p> <p>(A) $\frac{2}{25}$</p> <p>(B) $\frac{2}{5}$</p> <p>(C) $\frac{1}{2}$</p> <p>(D) $\frac{1}{32}$</p>	(D)
612	142	<p>A number of five digits is formed with digits 1, 2, 3, 4, 5 without repetition. The probability that it is a number divisible by 4 is</p> <p>(A) $\frac{1}{5}$</p> <p>(B) $\frac{2}{5}$</p> <p>(C) $\frac{3}{5}$</p> <p>(D) $\frac{4}{5}$</p>	(A)
612	143	<p>A bag contains 7 red and 5 white balls. Four white balls are drawn at random. The probability that all of them are red, is</p> <p>(A) $\frac{7}{99}$</p> <p>(B) $\frac{14}{33}$</p> <p>(C) $\frac{14}{99}$</p> <p>(D) $\frac{12}{35}$</p>	(A)
612	144	<p>A married couple appear for an interview for two vacancies in a company. The probability of man's selection is $\frac{3}{4}$ and that of woman's selection is $\frac{1}{3}$. The probability that both of them will be selected is</p> <p>(A) $\frac{1}{12}$</p> <p>(B) $\frac{5}{12}$</p>	(A)

		<p>(C) $\frac{1}{2}$</p> <p>(D) $\frac{7}{12}$</p>	
612	145	<p>In the group $G = \{1, 3, 7, 9\}$ under multiplication modulo 10, the inverse of 3 is</p> <p>(A) 1</p> <p>(B) 3</p> <p>(C) 7</p> <p>(D) 9</p>	(C)
612	146	<p>If G is a group and $a, b, c \in G$, then $(a b^{-1} c)^{-1}$ is</p> <p>(A) $a b c^{-1}$</p> <p>(B) $c^{-1} a^{-1} b$</p> <p>(C) $a^{-1} b c$</p> <p>(D) $c^{-1} b a^{-1}$</p>	(D)
612	147	<p>The number of improper subgroups of $G = \{1, -1, i, -i\}$ with respect to multiplication is</p> <p>(A) 2</p> <p>(B) 1</p> <p>(C) 3</p> <p>(D) 4</p>	(B)
612	148	<p>The sub-group $H = \{(1), (1\ 2)\}$ of S_3 is</p> <p>(A) an invariant subgroup of S_3</p> <p>(B) a normal subgroup of S_3</p> <p>(C) not a normal subgroup of S_3</p> <p>(D) a normal divisor of S_3</p>	(C)
612	149		(A)

		<p>From 6 men and 4 women , the number of ways of forming a committee of 5 members, if there is no restriction on its formation, is</p> <p>(A) 252</p> <p>(B) 240</p> <p>(C) 236</p> <p>(D) 180</p>	
612	150	<p>Let $a * b = a \times b - b$ ($a, b \in \mathbb{N}$ and $*$ is an operation on \mathbb{N}). Then $a * b = b * a$ implies</p> <p>(A) $a = 0$</p> <p>(B) $b = 0$</p> <p>(C) $a = b$</p> <p>(D) $a = -b$</p>	(C)