

CHEMISTRY PG





1. The toxic environmental pollutant dioxin contains 44.8% carbon, 1.25% hydrogen and 44.0% chlorine. Its molecular mass is  $320 \pm 10$ . What is the molecular formula of dioxin?
- (A)  $C_{10}H_8O_4Cl_4$   
(B)  $C_{12}H_6O_2Cl_4$   
(C)  $C_6H_2OCl_2$   
(D)  $C_{12}H_4O_2Cl_4$
2. Conversion of 2,5-dimethylfuran to 2,5-dimethylpyrrole (see equation) may be accomplished in two steps.
- (i) Hydrolytic ring opening of the furan in aqueous acid  
(ii) Heating the hydrolysis product with excess ammonium carbonate.

What is the intermediate hydrolysis product formed in first step that is subsequently used in the second step?




- (A) 2,5-hexanedione  
(B) 3,4-hexanedione  
(C) hexanedial  
(D) 2,5-dimethylcyclopentanone

3. A  $C_9H_{10}O$  compound has a strong absorption at 1730 and two smaller but sharp absorption peaks at 2719 and 2818  $cm^{-1}$  in the infrared.  $^1H$  NMR spectrum of the compound exhibited a 3 proton doublet at  $\delta 1.00$  ppm, a 1 proton doublet  $\delta 10.00$  ppm along with other multiplets. Which among the following compounds displays these features?




4. In the reaction of (*R*)-3-phenylbutan-2-one with methylmagnesium iodide to give 2-methyl-3-phenylbutan-2-ol, what happens to the configuration of the stereogenic center in it?




- (A) Nothing, it remains unchanged  
(B) Inversion takes place  
(C) Racemization occurs  
(D) The product is achiral

5. Which two of the following aldohexoses give the same osazone derivative?




- (A) 1 and 4  
 (B) 1 and 3  
 (C) 2 and 3  
 (D) 3 and 4





6. Assuming that their conformation is frozen as given below, which among the following  $\text{C}_8\text{H}_{16}$  isomers should have the lowest heat of combustion?







7. Which of the following trienes will have the longest wavelength absorption in its UV-Vis spectrum?



8. Which of the following is a D-aldopentose?




9. Which two of the following compounds are reduced to the same chiral alditol by sodium borohydride?



- (A) **1 and 2**  
 (B) **1, 2 and 4**  
 (C) **2 and 4**  
 (D) **3 and 4**

10. Which two of the following compounds, if any, are epimers?







- (A) **1** and **2**  
 (B) **3** and **4**  
 (C) **2** and **3**  
 (D) None of the above
11. Which among the following reagents is used in color test for amino acids?
- (A) ninhydrin  
 (B) cyanogen bromide  
 (C) trypsin  
 (D) chymotrypsin
12. Under identical reaction conditions which among the following compounds will undergo nitration at the fastest rate? (*Hint*: Baker Nathan Effect)




- (A) **I**  
 (B) **II**  
 (C) **III**  
 (D) **IV**
13. Which if any of the following compounds will display spin-spin splitting of protons in its  $^1\text{H}$  NMR spectrum?
- (A)  $(\text{CH}_3)_3\text{COCH}_3$   
 (B)  $\text{Br}(\text{CH}_2)_3\text{Br}$   
 (C) *p*-xylene (1,4-dimethylbenzene)  
 (D) None of the above

14. The  $^1\text{H}$  NMR spectrum of diethyl ether ( $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ ) shows
- (A) Two peaks, one a triplet, the other a quartet respectively in a 3:2 ratio
  - (B) Two peaks, one a triplet, the other a quartet respectively in a 2:3 ratio
  - (C) Four peaks, two 2 proton triplets and two 3 proton quartets
  - (D) Four peaks, two 3 proton triplets and two 2 proton quartets
15. Neopentyl chloride,  $(\text{CH}_3)_3\text{CCH}_2\text{Cl}$  ( $\text{C}_5\text{H}_9\text{Cl}$ ), reacts with the strong base sodium amide (can pick a C-H proton!) to form a new compound. This compound has a molecular ion at  $m/z = 70$  amu corresponding to the molecular formula  $\text{C}_5\text{H}_{10}$ , and displays two singlets at  $\delta$  0.20 and 1.05 ppm (intensity ratio = 2:3) in its  $^1\text{H}$  NMR spectrum. What is a plausible structure for this  $\text{C}_5\text{H}_{10}$  compound?
- (A) 2-methyl-2-butene
  - (B) 1,1-dimethylcyclopropane
  - (C) Methylcyclobutane
  - (D) Cyclopentane
16. Which of the following reactions **would not** be a useful method for preparing 1-phenylbutan-2-ol?
- (A) phenylacetaldehyde + ethylmagnesium bromide
  - (B) butanal + phenylmagnesium bromine
  - (C) propanal + benzylmagnesium bromine
  - (D) 1-phenylbutan-2-one +  $\text{NaBH}_4$
17. What can be the possible number of isomers of tribromobenzene?
- (A) 1
  - (B) 2
  - (C) 3
  - (D) 4

18. Among the four aromatic carboxylic acids given below, which one can be resolved into its enantiomers by reaction with an enantiomerically pure chiral amine?




19. How many stereoisomers do you expect for the following structural formula?




- (A) 2  
(B) 3  
(C) 4  
(D) 6

20. Arrange in order of C-H bond strength (C-H bond circled)




- (A) I > II > III
- (B) III > II > I
- (C) II > I > III
- (D) II > III > I

21. Which among these has highest acidity?



- (A) I
- (B) II
- (C) III
- (D) IV


22. The reagent/reagents used to convert I to II is



- (A)  $\text{PCl}_5$
- (B)  $\text{SOCl}_2$
- (C)  $\text{SOCl}_2 + \text{Pyridine}$
- (D)  $\text{HCl}$




23. Designate E/Z notation for the following:




- (A) I= E, II=E  
(B) I= Z, II=Z  
(C) I= E, II=Z  
(D) I= Z, II=E

24. Which among the following acids, forms anhydride fastest on heating?




25. Arrange the following carbanions in stability order (note: some of these carbanions may have more than one canonical form)







- (A) IV > I > II > III  
(B) II > I > IV > III  
(C) III > II > I > IV  
(D) I > IV > II > III

26. Arrange in the order of stability





- (A) I < II < III  
(B) II < I < III  
(C) I < III < II  
(D) II < III < I
27. Which of the following is a *meso* compound?
- (A) (2*R*,3*R*)-2,3-Dibromobutane  
(B) (2*R*,3*S*)-2,3-Dibromopentane  
(C) (2*R*,4*R*)-2,4-Dibromopentane  
(D) (2*R*,4*S*)-2,4-Dibromopentane
28. Ozonolysis of *cis*-but-2-ene followed by work-up and treatment with H<sub>2</sub>O<sub>2</sub> yields
- (A) CH<sub>3</sub>CHO  
(B) CH<sub>3</sub>COCH<sub>3</sub>  
(C) CH<sub>3</sub>CO<sub>2</sub>H  
(D) CH<sub>3</sub>CHO + CH<sub>3</sub>CO<sub>2</sub>H
29. Reactive intermediate involved in the conversion of bromobenzene to aniline in the presence of sodamide in liquid ammonia is
- (A) carbanion  
(B) free-radical  
(C) benzyne  
(D) carbocation
30. Grignard reagents react efficiently with carbon dioxide and various carbonyl compounds including acid halides. Products isolated after workup from the reaction of phenylmagnesium bromide with carbon dioxide and benzoyl chloride are
- (A) PhCO<sub>2</sub>H & Triphenyl carbinol (Ph<sub>3</sub>C-OH) respectively  
(B) PhCO<sub>2</sub>H (from both)  
(C) PhCHO ( from both)  
(D) PhCO<sub>2</sub>H & PhCOBr respectively

31. Acetaldehyde on reduction with  $\text{NaBH}_4$  yields product **M** which on treatment with acetyl chloride yields **N**. Two molecules of **N** underwent condensation in presence of sodium ethoxide and final work-up to yield **O**. The structure of **O** is



32. Benzoic acid can be converted to benzoyl chloride and ethyl benzoate in high yields using **X** and **Y** respectively. **X** and **Y** are



33. Absorbance of a solution showing 50% transmittance is


(A) 0.5

(B) 2


(C)  $\log 2$

(D)  $\log 0.5$





34. *N,N*-Dimethyl-2-aminobutane on successive treatment with excess methyl iodide followed by base yielded 1-butene as the product. Mechanism involved in the final elimination step is



35. Which among the following heteroaromatic compounds is more reactive towards nucleophiles?



- (A) I  
(B) III  
(C) IV  
(D) II
36. Acetone is treated with the Mannich reagent ( $\text{HCHO} + \text{Me}_2\text{NH} \cdot \text{HCl}$ ) to produce compound **X**. The latter is then heated to produce **Y**. The structures of **X** and **Y** are



37. The common feature of all alkaloids is that

- (A) All have aromatic ring residues  
(B) All have heterocyclic ring residues  
(C) All are polycyclic compounds  
(D) All contain N

38. Molecular formula (general) of a sesquiterpene is


- (A)  $\text{C}_{10}\text{H}_{16}$   
(B)  $\text{C}_5\text{H}_8$   
(C)  $\text{C}_{15}\text{H}_{24}$   
(D)  $\text{C}_{20}\text{H}_{32}$

39. Reaction of a carboxylic acid with one equivalent each of bromine and  $\text{PBr}_3$  followed by aqueous workup yields
- (A)  $\alpha$ -bromoacid
  - (B)  $\alpha$ -bromoester
  - (C) acyl bromide
  - (D) alkyl bromide
40. 1,3,5-Trihydroxybenzene is also called
- (A) Pyrogallol
  - (B) Phloroglucinol
  - (C) Quinol
  - (D) Resorcinol
41. Which among these is used to catalyze the conversion of propyne to acetone in presence of 40%  $\text{H}_2\text{SO}_4$ ?
- (A)  $\text{MgSO}_4$
  - (B)  $\text{CuSO}_4$
  - (C)  $\text{K}_2\text{SO}_4$
  - (D)  $\text{HgSO}_4$
42. Which oxidant is used successfully to convert toluene to benzaldehyde in high yields?
- (A)  $\text{KMnO}_4$
  - (B)  $\text{Cu}_2\text{O}$
  - (C)  $\text{CrO}_2\text{Cl}_2$
  - (D)  $\text{Cr}_2\text{O}_3$
43. Skraup synthesis of quinoline starts with
- (A) Glycol
  - (B) Glycerol
  - (C) Benzyl alcohol
  - (D) Amyl alcohol
44. Which is the chief source of morphine?
- (A) Tobacco
  - (B) Pepper
  - (C) Cinchona
  - (D) Poppy

45. Acetic acid on treatment with  $\text{PCl}_5$  yield **P** which on reaction with sodium azide yields **Q**. The latter on heating yielded **R**. Compounds **P**, **Q**, and **R** are


- (A)  $\text{CH}_3\text{Cl}$ ,  $\text{CH}_3\text{N}_3$ ,  $\text{CH}_3\text{NH}_2$
- (B)  $\text{CH}_3\text{COCl}$ ,  $\text{CH}_3\text{N}_3$ ,  $\text{CH}_3\text{NH}_2$
- (C)  $\text{CH}_3\text{COCl}$ ,  $\text{CH}_3\text{CON}_3$ ,  $\text{CH}_3\text{NCO}$
- (D)  $\text{CH}_3\text{Cl}$ ,  $\text{CH}_2\text{N}_2$ ,  $\text{CH}_4$

46. Among the following, how many obey Huckel's  $[4n+2]$  rule for aromaticity?



- (A) 4
- (B) 5
- (C) 6
- (D) 3

47. Identify the major product formed in the following reaction carried out under kinetically controlled conditions



- (A)
- (B)
- (C)
- (D)

48. L-DOPA is used for the treatment of
- (A) Tuberculosis
  - (B) Parkinson disease
  - (C) Diabetes
  - (D) Cancer
49. Among the following, the natural product which is not an alkaloid is
- (A) Reserpine
  - (B) Piperine
  - (C)  $\beta$ -Carotene
  - (D) Papavarine
50.  $n-\pi^*$  Transition in acetone will be most blue shifted in which among the following solvents?
- (A) Water
  - (B) Methanol
  - (C) Acetonitrane
  - (D) Hexane
51. Among the following molecules which one shows pure rotation spectra:
- (A)  $N_2$
  - (B)  $CO_2$
  - (C)  $H_2S$
  - (D)  $CH_4$
52. A particular vibration in a polyatomic molecule is IR active if during vibration, there is a change in
- (A) Polarizability
  - (B) Dipole moment
  - (C) Frequency
  - (D) Potential energy
53. Free radicals may be detected by
- (A) Mass spectrometry
  - (B) IR spectroscopy
  - (C) ESR spectroscopy
  - (D) UV- Visible spectroscopy

54. Match items in Column I with the appropriate items in Column II

Column I - Spectral Technique	Column II - Selection rule
(P) Rotational transition	(I) $\Delta v = \pm 1$
(Q) Vibrational Transition	(II) $\Delta J = \pm 1$
(R) Electronic Transition in atoms	(III) $\Delta v = 0$
	(IV) $\Delta l = \pm 1$
	(V) $\Delta l = 0$

- (A) (P)-(II), (Q)-(I), (R)-(IV)  
(B) (P)-(II), (Q)-(III), (R)-(V)  
(C) (P)-(III), (Q)-(II), (R)-(I)  
(D) (P)-(V), (Q)-(I), (R)-(II)

55. The packing fraction of a simple cubic lattice is close to

- (A) 0.94  
(B) 0.76  
(C) 0.52  
(D) 0.45

56. When Frenkel defects are created in an otherwise perfect ionic crystal, the density of the ionic crystal

- (A) increases  
(B) decreases  
(C) remains same  
(D) oscillates with the number of defects

57. A compound is formed by two elements M and N. The element N forms ccp lattice and atoms of M occupy  $\frac{1}{3}$ <sup>rd</sup> of tetrahedral voids. What is the formula of the compound?

- (A)  $MN_2$   
(B)  $M_2N_3$   
(C)  $M_3N_2$   
(D)  $M_2N_2$

58. A metal crystallizes in fcc structure with a unit cell side of 500 pm. If the density of the crystal is 1.33 g/cc, the molar mass of the metal is close to

- (A) 23  
(B) 24



- (C) 25  
(D) 26
59. A metal crystallizes in fcc lattice. The edge length of the unit cells is  $4.04 \times 10^{-8}$  cm and density is  $10.5 \text{ g cm}^{-3}$ . Calculate the atomic mass of the metal.
- (A) 144 g/mol  
(B) 125 g/mol  
(C) 106.6 g/mol  
(D) 213 g/mol
60. Which of the following statements is **not true** about the hexagonal close packing?
- (A) The coordination number is 12  
(B) It has 74% packing efficiency  
(C) Tetrahedral voids of the second layer are covered by the spheres of the third layer  
(D) In this arrangement spheres of the fourth layer are exactly aligned with those of the first layer
61. The standard cell potential for the reaction  $\text{Zn(s)} + \text{Cu}^{2+}(\text{aq}) \rightleftharpoons \text{Zn}^{2+}(\text{aq}) + \text{Cu(s)}$  is +1.10 V. The Gibbs free energy change during the reaction is ( $F = 96500 \text{ coulomb mol}^{-1}$ )
- (A)  $-21.2 \text{ kJ mol}^{-1}$   
(B)  $+212 \text{ kJ mol}^{-1}$   
(C)  $-212 \text{ kJ mol}^{-1}$   
(D)  $-212 \text{ J mol}^{-1}$
62. An absorbance shift towards a longer wavelength caused by chromophore is called
- (A) Hypochromic shift  
(B) Hypsochromic shift  
(C) Hyperchromic shift  
(D) Bathochromic shift
63. The rate of a chemical reaction doubles for every  $10^\circ\text{C}$  rise of temperature. If the temperature is raised by  $50^\circ\text{C}$ , the rate of the reaction increases by about
- (A) 32 times  
(B) 64 times  
(C) 10 times  
(D) 24 times

FOR REFERENCE ONLY

64. Choose the **incorrect** property of colloids
- (A) Colloidal particles do not settle down when left undisturbed that is colloidal sols are quite stable
  - (B) A colloidal solution is homogeneous in nature
  - (C) The size of colloidal particles lies in the range of  $10^{-5}$  -  $10^{-7}$  cm
  - (D) Colloidal particles are stabilized by surface charges
65. When  $O_2$  is adsorbed on a metallic surface, electron transfer occurs from the metal to  $O_2$ . The true statements regarding the adsorption are
- a.  $O_2$  is physisorbed
  - b. Heat is released
  - c. Occupancy of  $\pi^* 2p$  of  $O_2$  is increased
  - d. Bond length of  $O_2$  is increased
- (A) a, b, c
  - (B) b, c, d
  - (C) a, c, d
  - (D) a, b, d
66. Match the **Catalysts** to the correct **Processes**:
- | <b>Catalyst</b> | <b>Process</b>                    |
|-----------------|-----------------------------------|
| (P) $TiCl_3$    | (i) Wacker process                |
| (Q) $PdCl_2$    | (ii) Ziegler-Natta polymerisation |
| (R) $CuCl_2$    | (iii) Contact process             |
| (S) $V_2O_5$    | (iv) Deacon's process             |
- (A) (P)-(iii), (Q)-(ii), (R)-(iv), (S)-(i)
  - (B) (P)-(ii), (Q)-(i), (R)-(iv), (S)-(iii)
  - (C) (P)-(ii), (Q)-(iii), (R)-(iv), (S)-(i)
  - (D) (P)-(iii), (Q)-(i), (R)-(ii), (S)-(iv)
67. Rate of physisorption increases with
- (A) Decrease in temperature
  - (B) Increase in temperature at low temperatures
  - (C) Decrease in pressure
  - (D) Decrease in surface area
68. The process with negative entropy change is
- (A) Dissolution of iodine in water
  - (B) Sublimation of dry ice
  - (C) Synthesis of ammonia from  $N_2$  and  $H_2$
  - (D) Dissociation of  $CaSO_4(s)$  to  $CaO(s)$  and  $SO_3(g)$

69. An emulsion consists of
- (A) one liquid and one solid
  - (B) one liquid and one gas
  - (C) two liquids
  - (D) two solids
70. Which of the following statements regarding the second law of thermodynamics is true?
- (A) Energy cannot be created or destroyed, only transferred or transformed
  - (B) The total entropy of an isolated system can decrease over time
  - (C) Natural processes tend to move toward states of lower entropy
  - (D) Heat flows spontaneously from a hot body to a cold body
71. The heat given to an ideal gas in isothermal conditions is used to
- (A) Increase temperature
  - (B) Do external work
  - (C) Increase temperature and do external work
  - (D) Increase internal energy
72. What is the source temperature of the Carnot engine in K required to get 70% efficiency? Given: sink temperature = 27 °C
- (A) 1000 K
  - (B) 90 K
  - (C) 270 K
  - (D) 727 K
73. A gas undergoes a process in which its volume changes from  $V_1$  to  $V_2$  at constant pressure  $P$ . During this process,  $Q$  amount of heat is added to the gas, and  $W$  amount of work is done by the gas. If  $\Delta U$  represents the change in internal energy of the gas, which of the following equations correctly represents the relationship among  $Q$ ,  $W$ , and  $\Delta U$ ?
- (A)  $Q=W+P(V_2-V_1)$
  - (B)  $Q=\Delta U+W$
  - (C)  $\Delta U=Q+W$
  - (D)  $Q=W-\Delta U$

74. A PMR spectrometer operates at 300 MHz. Find the value of magnetic field.  
Given:  $g_N = 5.585$  and  $B_N = 5.05 \times 10^{-27} \text{ JT}^{-1}$
- (A) 7.05 T
  - (B) 6.38 T
  - (C) 7.58 T
  - (D) 5.93 T
75. Which of the following statement is **false** for mass spectroscopy?
- (A) Mass spectroscopy is used to identify unknown compounds within a sample, and to elucidate the structure and chemical properties of different molecules
  - (B) Particle are characterized by their mass to charge ratios ( $m/z$ ) and relative abundances
  - (C) This technique basically studies the effect of ionizing energy on molecules
  - (D) This technique can be used on all state of matter
76. Which spectroscopy is used for determination of heavy metal contamination in food?
- (A) Electrophoresis
  - (B) X-ray Diffraction
  - (C) ELISA
  - (D) Inductively coupled plasma spectroscopy
77. Which of the following statements about IR spectroscopy is **true**?
- (A) IR spectroscopy measures the absorption of visible light by a compound
  - (B) IR spectroscopy is commonly used to determine the molecular weight of a compound
  - (C) In IR spectroscopy, stretching vibrations typically occur at higher wave numbers than bending vibrations
  - (D) IR spectroscopy can distinguish between cis and trans isomers based on their absorption spectra
78. Choose the correct sequence of process during Atomization in atomic absorption spectroscopy
- (A) Desolvation – Nebulization – Dissociation – Volatilization – Ionization
  - (B) Nebulization – Desolvation – Volatilization – Dissociation – Ionization
  - (C) Desolvation – Nebulization – Volatilization – Dissociation – Ionization
  - (D) Nebulization – Volatilization – Desolvation – Dissociation – Ionization

79. What is the heat of formation of ethane?

(Given that the heat of combustion of ethane ( $C_2H_6$ ), hydrogen ( $H_2$ ) and carbon (C) are  $-1411$  kJ/mol,  $-393$  kJ/mol,  $-286$  kJ/mol respectively.)

- (A)  $-240$  kJ/mol
- (B)  $240$  kJ/mol
- (C)  $-340$  kJ/mol
- (D)  $440$  kJ/mol

80. Which of the following is true for the process of spontaneous adsorption of a gaseous component?

- (A)  $\Delta G < 0, \Delta S < 0, \Delta H > 0$
- (B)  $\Delta G < 0, \Delta H < 0, \Delta S > 0$
- (C)  $\Delta H > 0, \Delta S > 0, \Delta G < 0$
- (D)  $\Delta H < 0, \Delta G < 0, \Delta S < 0$

81. Which of the following is Langmuir adsorption isotherm?

(NOTE:  $x$  = amount of substance adsorbed,  $m$  = mass of adsorbent,  $p$  = pressure,  $a$  and  $b$  = constants)

(A)  $\frac{m}{x} = \frac{1+bp}{ap}$

(B)  $\frac{x}{m} = \frac{ap}{1+bp}$

(C)  $\frac{x}{m} = \frac{ap}{1+ap}$

(D)  $\frac{m}{x} = \frac{ap}{1+bp}$

82. The isotope  $^{90}\text{Sr}$  emits radiation by a first order process and has a half-life of 23 years. When ingested by humans, it becomes permanently incorporated in bone tissue. If  $1 \mu\text{g}$  is administered to a person, how much of this isotope remains in the body after 34 years?


- (A)  $0.86 \mu\text{g}$
- (B)  $0.36 \mu\text{g}$
- (C)  $0.61 \mu\text{g}$
- (D)  $0.55 \mu\text{g}$

83. For the reaction  $A + B \rightarrow \text{products}$ , it is observed that
- On doubling the concentration of B only, the rate of reaction is also doubled.
  - On doubling the initial concentration of both A and B, there is change by a factor of 8 in the rate of reaction.

The rate of reaction is given by

- rate =  $k [A]^2 [B]$
- rate =  $k [A] [B]^2$
- rate =  $k [A]^2 [B]^2$
- rate =  $k [A] [B]$

84. Which of the following plots yields a straight line graph for a second order reaction?



$[A]_0$  = initial concentration of A;  $[A]_t$  = concentration of A at time = t

- Half-life vs.  $1/[A]_0$
- Half-life vs.  $[A]_0$
- Half-life vs.  $\ln[A]_0$
- Half-life vs.  $[A]_t$


85. The rise of liquid in a capillary does **NOT** depend upon

- Angle of contact
- Radius of the capillary
- Atmospheric pressure
- Density of the liquid

86. Calculate the ionic strength in a solution that contains 0.0100 mol KCl, 0.0050 mol  $MgCl_2$ , 0.0020 mol  $MgSO_4$  and 100 g  $H_2O$ .

- 0.68
- 0.17
- 0.33
- 0.19

87. Calculate  $E_{\text{cell}}$  for the given cell at  $25^\circ\text{C}$  and 1 bar given that activity coefficient of  $\text{CuSO}_4$  is 0.043:



(NOTE: Standard reduction potential for the  $\text{Cu}|\text{CuSO}_4$  electrode is 0.339 V and for the  $\text{Hg}|\text{Hg}_2\text{SO}_4(\text{c})$  electrode is 0.615 V at 298 K.)

- (A) 0.122 V  
(B) 0.243 V  
(C) 0.542 V  
(D) 0.316 V
88. For a given amount of an ideal gas the volume was increased by four times and the temperature was increased by two times. What will be the change in pressure of the gas?
- (A) Increases 2 times  
(B) Increases 4 times  
(C) Remains same  
(D) Reduces to half
89. The area under the curve  $\int TdS$  is equal to
- (A) Change in entropy  
(B) Work done in reversible conditions  
(C) Total heat transferred reversibly  
(D) Change in internal energy
90. Of the following, which will effectively lower the activation energy?
- (A) Increasing concentrations of reactants  
(B) Adding a catalyst for the reaction  
(C) Raising the temperature of the reaction  
(D) Removing products as reaction proceeds



91. The reaction  $A + B \rightarrow P$  is endothermic and has an activation energy of 75 kJ/mol. Which of the following statements are correct?
- I. The reverse reaction has an activation energy equal to 75 kJ/mol.
  - II. The reverse reaction has an activation energy greater than 75 kJ/mol.
  - III. The reverse reaction has an activation energy less than 75 kJ/mol.
  - IV. The change in internal energy is greater than zero.
  - V. The change in internal energy is less than zero.
- (A) I and IV  
(B) II and V  
(C) III and V  
(D) III and IV
92. Which of the following is the correct set of quantum numbers for outermost electron of potassium ( $Z = 19$ )?
- (A)  $n = 4, l = 3, m = 2, s = -\frac{1}{2}$   
(B)  $n = 4, l = 2, m = 0, s = -\frac{1}{2}$   
(C)  $n = 4, l = 1, m = 0, s = \frac{1}{2}$   
(D)  $n = 4, l = 0, m = 0, s = -\frac{1}{2}$
93. If the electronic configuration of nitrogen ( $Z = 7$ ) is written as  $(1s^2, 2s^2, 2p_x^2, 2p_y^1)$ , which of the following rules would be violated?
- (A) Pauli exclusion principle  
(B) Hund's rule of maximum multiplicity  
(C) Aufbau's principle  
(D) Bohr's atomic rule
94. The relation  $\Delta x \cdot \Delta p = \frac{h}{4\pi}$  represents
- (Note: symbols 'x' and 'p' carry their conventional meaning in quantum mechanics)
- (A) Heisenberg's uncertainty principle  
(B) de Broglie equation  
(C) Pauli exclusion principle  
(D) Schrodinger wave equation
95. In  $dsp^2$  hybridisation, what is the general geometry of the newly formed molecular orbitals expected to be?
- (A) Tetrahedral  
(B) Square planar  
(C) Trigonal pyramidal  
(D) Trigonal bipyramidal

96. The solubility of a salt is 's' and its solubility product is '4s<sup>3</sup>'. What is the ratio of cations and anions in the salt?
- (A) 1 : 2  
(B) 1 : 1  
(C) 2 : 2  
(D) 1 : 3
97. Let  $\lambda$  be the equivalent conductance of an electrolyte at a given concentration and  $\lambda_{\infty}$  be the equivalent conductance at infinite dilution. What would be the correct expression for degree of dissociation ( $\alpha$ ) of the electrolyte?
- (A)  $\alpha = \lambda_{\infty} - \lambda$   
(B)  $\alpha = \lambda_{\infty} / \lambda$   
(C)  $\alpha = \lambda - \lambda_{\infty}$   
(D)  $\alpha = \lambda / \lambda_{\infty}$
98. The equilibrium constant for the reaction  $3\text{C}_2\text{H}_2 \rightleftharpoons \text{C}_6\text{H}_6$  at 298 K is 4. If the equilibrium concentration of  $\text{C}_2\text{H}_2$  is 0.5 mol/L, what would be the concentration of  $\text{C}_6\text{H}_6$  at that point?
- (A) 1.2 M  
(B) 0.1 M  
(C) 2.3 M  
(D) 0.5 M
99. The function  $\Phi = e^{ax}$  is an eigenfunction of the operator  $\left(\frac{d^2}{dx^2}\right)$ . What would be the resulting eigenvalue?
- (A)  $2a$   
(B)  $a^2$   
(C)  $a$   
(D)  $2a^2$
100. The Daniel cell is
- (A)  $\text{Zn}_{(s)} | \text{Zn}^{2+}_{(aq)} || \text{Cu}^{2+}_{(aq)} | \text{Cu}_{(s)}$   
(B)  $\text{Zn}_{(s)} | \text{Zn}^{2+}_{(aq)} || \text{Ag}^{2+}_{(aq)} | \text{Ag}_{(s)}$   
(C)  $\text{Fe}_{(s)} | \text{Fe}^{2+}_{(aq)} || \text{Cu}^{2+}_{(aq)} | \text{Cu}_{(s)}$   
(D)  $\text{H}_{2(s)} | \text{H}_2\text{SO}_{4(aq)} || \text{Cu}^{2+}_{(aq)} | \text{Cu}_{(s)}$

101. The bond orders in  $N_2$  and  $[N_2]^-$  are
- (A) 3 and 2.5
  - (B) 3 and 3
  - (C) 2.5 and 3
  - (D) 2.5 and 2.5
102. Identify the geometry of the following species
- I =  $:N(SiH_3)_3$    II =  $:N(CH_3)_3$    III =  $\bullet CF_3$    IV =  $\bullet CH_3$
- (A) I & II pyramidal, III & IV planar
  - (B) I & III pyramidal, II & IV planar
  - (C) I, II & III pyramidal, IV is planar
  - (D) I & IV planar, II & III pyramidal
103. The number of metal-metal bonds present in  $Co_4(CO)_{12}$  is
- (A) 4
  - (B) 5
  - (C) 6
  - (D) 7
104. Among  $Si(CH_3)_4$ ,  $NCl_3$ ,  $trans-[SnCl_4(py)_2]$  ( $py$  = pyridine),  $P(O)Cl_3$ , those with zero dipole moment are
- (A)  $Si(CH_3)_4$  &  $NCl_3$
  - (B)  $POCl_3$  &  $trans-[SnCl_4(py)_2]$
  - (C)  $Si(CH_3)_4$  &  $trans-[SnCl_4(py)_2]$
  - (D)  $POCl_3$  &  $Si(CH_3)_4$
105. The half-life of the first excited state of  $^{57}Fe$  is  $1.5 \times 10^{-7}$  s. The line width of the resonance is
- (A) 4.85 J
  - (B)  $4.86 \times 10^{-28}$  J
  - (C) 4.55 J
  - (D)  $4.58 \times 10^{-20}$  J

106. The correct order of second ionization energy ( $IE_2$ ) values of C, N, O and F is
- (A)  $O > F > N > C$
  - (B)  $F > O > N > C$
  - (C)  $F > O > C > N$
  - (D)  $N > C > F > O$
107. How many signals will you observe in the ESR spectrum of  $^2\text{H}$ -atom?
- (A) 2
  - (B) 4
  - (C) 3
  - (D) 1
108. Which of the following pair is not iso-structural?
- (A)  $\text{XeO}_4$  &  $\text{S}_2\text{O}_3^{2-}$
  - (B)  $\bullet\text{CH}_3$  (radical) &  $\bullet\text{CF}_3$  (radical)
  - (C)  $\text{SOCl}_2$  &  $\text{NH}_3$
  - (D)  $\text{ICl}_4^-$  &  $\text{XeF}_4$
109. Unit less property among the following is
- (A) Electron potential
  - (B) Electron affinity
  - (C) Ionization enthalpy
  - (D) Electronegativity
110. When NaCl crystal is heated in Na vapour, it develops a yellow colour, owing to trapping of an electron in an anion vacancy. The colour centre is called
- (A) M - centre
  - (B) R- centre
  - (C)  $F_A$ - centre
  - (D) F- centre
111. The increasing order of Brönsted acidity of boron hydrides is
- (A)  $\text{B}_5\text{H}_9 < \text{B}_4\text{H}_{10} < \text{B}_{10}\text{H}_{14}$
  - (B)  $\text{B}_4\text{H}_{10} < \text{B}_{10}\text{H}_{14} < \text{B}_5\text{H}_9$
  - (C)  $\text{B}_4\text{H}_{10} < \text{B}_5\text{H}_9 < \text{B}_{10}\text{H}_{14}$
  - (D)  $\text{B}_{10}\text{H}_{14} < \text{B}_5\text{H}_9 < \text{B}_4\text{H}_{10}$

112. An insulator NiO when heated in air to 1000 °C produce a black colored defect NiO which behaves as a semiconductor. The semiconductor is
- (A) Metal excess semiconductor
  - (B) Controlled valency semiconductor
  - (C) p-type
  - (D) n-type
113. The structure of O<sub>3</sub> and N<sub>3</sub><sup>-</sup> are
- (A) Linear and bent respectively
  - (B) Both linear
  - (C) Both bent
  - (D) Bent and linear respectively
114. Which of the following is the correct meaning of an improper rotational axis of symmetry?
- (A) Rotation followed by reflection
  - (B) Rotation followed by translation
  - (C) Rotation by 180°
  - (D) Reflection followed by translation
115. Which of the following is the ground state spectroscopic term symbol for free V<sup>3+</sup> ion?
- (A) <sup>3</sup>F
  - (B) <sup>3</sup>P
  - (C) <sup>4</sup>F
  - (D) <sup>4</sup>P
116. In Mn<sub>3</sub>O<sub>4</sub>, the number of manganese ions in tetrahedral and octahedral sites are respectively
- (A) One Mn<sup>3+</sup> and two Mn<sup>2+</sup>
  - (B) One Mn<sup>2+</sup> and two Mn<sup>3+</sup>
  - (C) Two Mn<sup>3+</sup> and one Mn<sup>2+</sup>
  - (D) Two Mn<sup>2+</sup> and one Mn<sup>3+</sup>

117. Which of the following is responsible for higher  $\mu_{\text{eff}}$  value of  $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$  than that of  $[\text{CoCl}_4]^{2-}$ ?
- (A) spin-orbit coupling in  $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$
  - (B) spin-orbit coupling in  $[\text{CoCl}_4]^{2-}$
  - (C) orbital contribution in  $[\text{CoCl}_4]^{2-}$
  - (D) orbital contribution in  $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$
118. A  $\text{Cr}^{3+}$  metal complex has strong transitions at  $\lambda_{\text{max}}$  at 431 nm, 781 nm, and 1,250 nm in an octahedral field. What will be the  $\Delta_{\text{oct}}$  for this complex?
- (A) 23,200  $\text{cm}^{-1}$
  - (B) 12,805  $\text{cm}^{-1}$
  - (C) 8,000  $\text{cm}^{-1}$
  - (D) 4,805  $\text{cm}^{-1}$
119. What kind of indicator is Ferroin?
- (A) acid-base indicator
  - (B) complexometric indicator
  - (C) pH indicator
  - (D) redox indicator
120. A transition metal forms two octahedral complexes in the same oxidation state. One of them is having a magnetic moment 4.9 B.M., while that of another is 0.0 B.M. The metal ion that fit this description is
- (A)  $\text{Cr}(\text{II})$
  - (B)  $\text{Mn}(\text{II})$
  - (C)  $\text{Fe}(\text{II})$
  - (D)  $\text{Ni}(\text{II})$
121. The complex which follows the 18-electron rule is
- (A)  $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2]$
  - (B)  $[(\eta^5\text{-C}_5\text{H}_5)\text{Mo}(\text{CO})_3]$
  - (C)  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Co}]$
  - (D)  $[(\eta^5\text{-C}_5\text{H}_5)\text{Re}(\eta^6\text{-C}_6\text{H}_6)]$

122. The orbital hybridization involved in  $\text{CrO}_4^{2-}$  is
- (A)  $\text{sp}^3$
  - (B)  $\text{sd}^3$
  - (C)  $\text{spd}^2$
  - (D)  $\text{sp}^2\text{d}$
123. The point group of  $\text{ClF}_3$  is
- (A)  $\text{C}_{2v}$
  - (B)  $\text{C}_s$
  - (C)  $\text{D}_{3h}$
  - (D)  $\text{C}_{3v}$
124. The correct order of electron transfer rate in the reaction is
- $$[\text{Co}(\text{NH}_3)_5\text{X}]^{2+} + [\text{Cr}(\text{OH}_2)_6]^{2+} \rightarrow [\text{Co}(\text{OH}_2)_6]^{3+} + [\text{Cr}(\text{OH}_2)_5\text{X}]^{2+} + 5\text{NH}_3$$
- $\text{X} = \text{F}^-, \text{Cl}^-, \text{Br}^-$  and  $\text{I}^-$
- (A)  $\text{I}^- < \text{F}^- < \text{Cl}^- < \text{Br}^-$
  - (B)  $\text{F}^- > \text{Cl}^- > \text{Br}^- > \text{I}^-$
  - (C)  $\text{F}^- < \text{Cl}^- > \text{Br}^- < \text{I}^-$
  - (D)  $\text{F}^- < \text{Cl}^- < \text{Br}^- < \text{I}^-$
125. Which of the following doesn't contribute to crystal field splitting parameter  $10Dq$ ?
- (A) Nuclear charge
  - (B) Spin-orbit coupling
  - (C) Electron and neutron interaction
  - (D) Electron and nucleus distance
126. Which of the following is considered as cellular storehouse of iron?
- (A) Ceruloplasmin
  - (B) Transferrin
  - (C) Ferroportin
  - (D) Ferritin
127. Kroll process is adopted for the extraction of
- (A) Sc
  - (B) Zr
  - (C) V
  - (D) Hf

128. Which of the following is meant for, the orbitals on the metal ion are smeared out over a larger region of space whenever the ligands bind to a metal ion?
- (A) j-j coupling effect
  - (B) nephelauxetic effect
  - (C) lanthanide contraction effect
  - (D) LS coupling effect
129. Which of the following is **NOT** a reducing agent?
- (A)  $\text{NaBH}_4$
  - (B)  $\text{LiAlH}_4$
  - (C)  $\text{NH}_2\text{NH}_2$
  - (D)  $\text{Na}_2\text{S}_2\text{O}_8$
130. Which of the following are the products formed when  $\text{XeF}_4$  reacts with water
- (A)  $\text{XeOF}_2 + 2\text{HF}_2$
  - (B)  $\text{XeO}_3 + \text{Xe} + \text{HF} + \text{O}_2$
  - (C)  $\text{XeOF}_4 + \text{XeF}_2 + \text{HF}$
  - (D)  $\text{XeO}_3 + \text{HF}$
131. The structure of  $\text{Ni}(\text{CO})_4$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  complexes are
- (A) both square planar
  - (B) square planar and tetrahedral respectively
  - (C) tetrahedral and square planar respectively
  - (D) both tetrahedral
132. The correct electronic configuration and magnetic moment of  $\text{Gd}^{3+}$  is
- (A)  $[\text{Xe}]4f^7$  and 7.9 B.M.
  - (B)  $[\text{Xe}]5f^7$  and 7.9 B.M.
  - (C)  $[\text{Xe}]4f^6$  and 8.9 B.M.
  - (D)  $[\text{Xe}]4f^9$  and 11.9 B.M.
133. The spin-only magnetic moment (in B.M.) value of  $[\text{Fe}(\text{CN})_6]^{3-}$  ion is
- (A) 5.916
  - (B) 1.73
  - (C) 0
  - (D) 1.414



134. Predict the bond order variation in the following  $O_2, O_2^+, O_2^-, O_2^{2-}$
- (A)  $O_2^+ > O_2 > O_2^- > O_2^{2-}$   
(B)  $O_2^{2-} > O_2^- > O_2 > O_2^+$   
(C)  $O_2^{2-} > O_2 > O_2^- > O_2^+$   
(D)  $O_2^{2-} > O_2^- > O_2^+ > O_2$
135. The correct order of electron gain enthalpies of P, S, Cl and F is
- (A)  $F > Cl > S > P$   
(B)  $Cl > F > P > S$   
(C)  $F > Cl > P > S$   
(D)  $Cl > F > S > P$
136. Three centre two electron bond is present in
- (A)  $BF_3$   
(B)  $B_2H_6$   
(C)  $H_3BO_3$   
(D)  $AlCl_3$
137. Which among the following doesn't obey 18-electron rule?
- (A)  $V(CO)_6$   
(B)  $Mn_2(CO)_{10}$   
(C)  $Fe(CO)_5$   
(D)  $Ni(CO)_4$
138. Among the oxides of nitrogen,  $N_2O_3, N_2O_4, N_2O_5$  the compound(s) having N–N bond(s) is/are
- (A)  $N_2O_4$  and  $N_2O_5$   
(B)  $N_2O_3$  and  $N_2O_5$   
(C)  $N_2O_4$  and  $N_2O_3$   
(D)  $N_2O_5$  only

139. Which of the following is not considered as an organometallic compound?
- (A) Ferrocene
  - (B) Cis-platin
  - (C) Ziese's salt
  - (D) Grignard reagent
140. A stable oxidation state of +4 is shown by
- (A) La
  - (B) Yb
  - (C) Sm
  - (D) Ce
141. Which of the following does not exist?
- (A) XeOF<sub>4</sub>
  - (B) XeF<sub>4</sub>
  - (C) ArF<sub>3</sub>
  - (D) XePtF<sub>6</sub>
142. The indicator used in the complexometric titration of ZnSO<sub>4</sub> and EDTA is
- (A) Diphenylamine
  - (B) N-Phenylanthranilic acid
  - (C) Thymol blue
  - (D) Eriochrome black -T
143. The crystal field stabilization energy of [Fe(CN)<sub>6</sub>]<sup>2-</sup> is ( $\Delta_0$  is the splitting energy and P is the pairing energy).
- (A)  $2.4 \Delta_0 - 3P$
  - (B)  $1.2 \Delta_0 - 3P$
  - (C)  $2.4 \Delta_0 - P$
  - (D) 0
144. Dimethylglyoxime (DMG) is used in spot test for confirming which cation?
- (A) Ni<sup>2+</sup>
  - (B) Zn<sup>2+</sup>
  - (C) K<sup>+</sup>
  - (D) Pt<sup>4+</sup>

145. Which of the following complex ions is diamagnetic in nature?
- (A)  $[\text{CoF}_3]^{3-}$
  - (B)  $[\text{Ni}(\text{CN})_4]^{2-}$
  - (C)  $[\text{NiCl}_4]^{2-}$
  - (D)  $[\text{Fe}(\text{CN})_4]^-$
146. The complexes,  $[\text{Co}(\text{NH}_3)_3][\text{Cr}(\text{CN})_6]$  and  $[\text{Cr}(\text{NH}_3)_3][\text{Co}(\text{CN})_6]$  are the examples of
- (A) Geometrical isomerism
  - (B) Linkage isomerism
  - (C) Ionisation isomerism
  - (D) Coordination isomerism
147. The magnetic moment of a transition metal ion is  $\sqrt{15}$  BM. Therefore the number of unpaired electrons present in it is
- (A) 1
  - (B) 2
  - (C) 3
  - (D) 4
148. Name the type of structure of silicate in which only one oxygen atom of  $[\text{SiO}_4]^{4-}$  is shared.
- (A) Three dimensional silicate
  - (B) Ambhibole
  - (C) Sheet silicate
  - (D) Pyrosilicate
149. In which of the following set is the bond order 3 for all species?
- (A) NO,  $\text{N}_2$  and CO
  - (B) NO,  $\text{N}_2$ ,  $\text{CN}^-$
  - (C)  $\text{NO}^+$ ,  $\text{N}_2$ , CO
  - (D)  $\text{NO}^+$ ,  $\text{N}_2$ ,  $\text{CN}^-$

150. In the reaction  $2X + B_2H_6 \rightarrow [BH_2(X)_2]^+[BH_4]^-$ , the amine X is

- (A)  $(CH_3)_3N$
- (B)  $(CH_3)_2NH$
- (C)  $CH_3NH_2$
- (D)  $NH_3$

FOR REFERENCE ONLY

## ANSWER KEY

**Subject Name: 604 CHEMISTRY**

SI No.	Key	SI No.	Key	SI No.	Key	SI No.	Key	SI No.	Key
1	D	31	B	61	C	91	D	121	D
2	A	32	B	62	D	92	D	122	B
3	C	33	C	63	A	93	B	123	A
4	A	34	C	64	B	94	A	124	D
5	B	35	C	65	B	95	B	125	C
6	C	36	B	66	B	96	A	126	D
7	D	37	D	67	A	97	D	127	B
8	B	38	C	68	C	98	D	128	B
9	A	39	A	69	C	99	B	129	D
10	C	40	B	70	D	100	A	130	B
11	A	41	D	71	B	101	A	131	C
12	A	42	C	72	A	102	D	132	A
13	B	43	B	73	B	103	C	133	B
14	A	44	D	74	A	104	C	134	A
15	B	45	C	75	D	105	B	135	D
16	B	46	A	76	D	106	A	136	B
17	C	47	A	77	C	107	C	137	A
18	C	48	B	78	C	108	B	138	C
19	D	49	C	79	C	109	D	139	B
20	B	50	A	80	D	110	D	140	D
21	A	51	C	81	C	111	C	141	C
22	B	52	B	82	B	112	C	142	D
23	C	53	C	83	A	113	D	143	A
24	A	54	A	84	A	114	A	144	A
25	D	55	C	85	C	115	A	145	B
26	D	56	C	86	C	116	B	146	D
27	D	57	B	87	D	117	D	147	C
28	C	58	C	88	D	118	C	148	D
29	C	59	C	89	C	119	D	149	C
30	A	60	D	90	B	120	C	150	D